

Administrarea subcutanata a medicamentelor in ingrijirea paliativa



Luciana Negulescu
Asistent medical
Asociatia pentru Servicii Mobile de Ingrijire Paliativa
Bucuresti

Administrarea subcutanata a medicamentelor

Indicatii:

- tumori sfera ORL,
- varsaturi,
- disfagie,
- stare terminala,
- coma.



Administrarea subcutanata a medicamentelor

Avantaje:

- evitarea administrării IV, IM (durere, spargerea venei)
- frecventa redusa a intepaturilor,
- bun control simptomatic,
- fixarea acului (functional 5-7 zile),
- confort crescut pentru pacient inclusiv la domiciliu.



Administrarea subcutanata a medicamentelor

Dezavantaje:

- nu toate medicamentele pot fi administrate SC (substante uleioase sau iritante),
- maxim 2-2,5 ml la o administrare.

Incidente (rare):

- posibila durere la locul injectiei,
- sangerare,
- infundarea acului,
- iritatie locala*.



Administrarea subcutanata a medicamentelor

Medicamentele care pot fi administrate SC:

Antialgice: ketorolac
tramadol
morfina



Administrarea subcutanata a medicamentelor

Medicamentele care pot fi administrate SC:

Antiemetice: metoclopramid
ondansetron
scobutil



Administrarea subcutanata a medicamentelor

Medicamentele care pot fi administrate SC:

Altele:

dexametazona

haloperidol

midazolam

levomepromazina

fenobarbital


ranitidina

atropina



Administrarea subcutanata a medicamentelor

Zone ale administrarii SC

















































Administrarea subcutanata a medicamentelor

Materiale folosite la administrarea SC:

- microperfuzoare (fluturasi) cu dimensiunea de 25 G,
- siringa 2 ml,
- pansament ocluziv transparent “tegaderm”, pentru fixare ac
- tampon cu alcool.



Compatibilitatea substantelor in siringa

	Morfina	Midazo -lam	Haloperi -dol	Scobu -til	Dexame -tazona	Metoclo -pramid	Trama -dol	Raniti -dina
Morfina							—	
Midazo -lam								
Haloperi -dol								?
Scobutil								
Dexame -tazona								?
Metoclo -pramid								?
Trama -dol	—							?
Raniti -dina			?		?	?	?	

Bibliografie

Ferrel B, Coyle N, Textbook of Palliative Nursing, 2nd Edition Oxford University Press, 2007

Negro S, Azuara ML, Sanchez Y, Reyes R, Barcia E, *Physical compatibility and in vivo evaluation of drug mixture for subcutaneous infusion to cancer patients in palliative care*, Supportive Care in Cancer, 2002

Neuenschwander H et al, *Medicine palliative*, Ligue suisse contre le cancer, 2007

